

r -CLEAN RINGS

NAHID ASHRAFI AND EBRAHIM NASIBI

ABSTRACT. An element of a ring R is called clean if it is the sum of an idempotent and a unit. A ring R is called clean if each of its element is clean. An element $r \in R$ called regular if $r = ryr$ for some $y \in R$. The ring R is regular if each of its element is regular. In this paper we define a ring is r -clean if each of its elements is the sum of a regular and an idempotent element. We give some relations between r -clean and clean rings. Finally we investigate some properties of r -clean rings.

1. INTRODUCTION

Throughout this paper, R denotes an associative ring with identity, $U(R)$ the group of units, $Id(R)$ the set of idempotents, $J(R)$ the Jacobson radical and $M_n(R)$ the ring of all $n \times n$ matrices over R .

An element x of a ring is called clean if $x = u + e$, where $u \in U(R)$ and $e \in Id(R)$. A ring R is called clean if each of its element is clean. Clean rings first were introduced by Nicholson [9]. Several peoples worked on this subject and investigate properties of clean rings, for example see [1], [3], [6] and [11].

A ring R is said to be exchange ring if for each $a \in R$ there exists $e \in Id(R)$ such that $e \in aR$ and $(1 - e) \in (1 - a)R$. Nicholson [9, Proposition 1.8] proved that clean rings are exchange and a ring with central idempotents is clean if and only if it is exchange. Camillo and Yu [3, Theorem 9] proved that a ring is semiperfect if and only if it is clean and has no infinite orthogonal family of idempotents.

In 1936, von Neumann defined that an element $r \in R$ is regular if $r = ryr$ for some $y \in R$, the ring R is regular if each of its element is regular. Some properties of regular rings has been studied in [5]. A ring R is called unit regular if, for each $a \in R$, there exists a unit $u \in R$ such that $aua = a$. Camillo and Yu [3, Theorem 5] proved that every unit regular ring is clean. In [10] Nicholson and Varadarajan proved that the converse is not true.

Let $Reg(R) = \{a \in R : a \text{ is regular}\}$. We call an element x of a ring R is r -clean if $x = r + e$, where $r \in Reg(R)$ and $e \in Id(R)$. A ring R is r -clean if each of its element is r -clean.

Clearly regular rings and clean rings are r -clean. But we will give some examples that shows in general, r -clean rings may not be regular. Also we will give an example that shows in general, r -clean rings may not be clean. In fact Bergman [7, Example 1] has constructed a regular ring which is not directly finite (a ring R is directly finite if for any elements $a, b \in R$, $ab = 1$

2000 *Mathematics Subject Classification.* 16E50, 16U99.

Key words and phrases. Bergman's example, clean rings, r -clean rings, von Neumann regular rings.

implies $ba = 1$) with 2 invertible in which not every element is a sum of units. We will prove that Bergman's example is not clean, but clearly it is r -clean.

We will show that a directly finite ring $R \neq 0$ is local if and only if it is r -clean and 0 and 1 are the only idempotents in R . Finally we give some properties of r -clean rings and we will prove that if R is an r -clean ring, then so is the matrix ring $M_n(R)$ for any $n \geq 1$.

2. r -CLEAN RINGS

In this section first we define r -clean element and r -clean rings and we show that the class of clean rings are a proper subset the class of r -clean rings.

Definition 1. *An element x of a ring R is r -clean if $x = r + e$, where $r \in \text{Reg}(R)$ and $e \in \text{Id}(R)$, a ring R is r -clean if each of its element is r -clean.*

Clearly regular rings and clean rings are r -clean. But in general, r -clean rings may not be regular. For example, every semiperfect ring is clean (see [3, Theorem 9]), so it is r -clean. But clearly it is not regular. Further $\mathbb{Z}_4$ is not regular, because $\bar{2}$ is not regular in $\mathbb{Z}_4$, but it is easy to check that $\mathbb{Z}_4$ is r -clean.

Also in general, r -clean rings may not be clean. For example [7, Example 1], proceeding as Bergman's example, let F be a field with $\text{char}(F) \neq 2$, $A = F[[x]]$ and K be the field of fractions of A . All the ideals of A are generated by power of x , denote by (x^n) . Define:

$$R = \{r \in \text{End}(A_F) : \text{there exists } q \in K \text{ and a positive integer } n, \\ \text{with } r(a) = qa \text{ for all } a \in (x^n)\}.$$

By [7, Example 1], R is a regular ring which is not directly finite and R is not generated by its units. So every element of R is not a sum of units, and since $\text{char}(F) \neq 2$, 2 is invertible in R . Also since R is regular thus R is r -clean. But R is not clean, because in otherwise, Proposition 10 in [3] (Let R be a ring in which 2 is invertible. Then R is clean if and only if every element of R is the sum of a unit and a square root of 1) implies that every element in R is a sum of a unit and a square root of 1. Thus every element in R is a sum of two units which it is a contradiction.

In following, we investigate some conditions in which r -clean rings are clean.

Lemma 2. *A ring $R \neq 0$ is local if and only if it is clean and 0 and 1 are the only idempotents in R .*

Proof. See [11, Lemma 14]. □

Theorem 3. *If $R \neq 0$ is a directly finite r -clean ring and 0 and 1 are the only idempotents in R , then R is clean.*

Proof. Since R is r -clean, each $x \in R$ has the form $x = r + e$, where $r \in \text{Reg}(R)$ and $e \in \text{Id}(R)$. If $r = 0$, then $x = e = (2e - 1) + (1 - e)$ and since $2e - 1 \in U(R)$ and $1 - e \in \text{Id}(R)$, so x is clean. Hence R is clean. But if $r \neq 0$, then there exists $y \in R$ such that $ryr = r$. Thus $ry \in \text{Id}(R)$. So by hypothesis, $ry = 0$ or $ry = 1$. Now if $ry = 0$, then $r = ryr = 0$ which is a

contradiction. Therefore $ry = 1$ and since R is directly finite so $yr = ry = 1$. Thus $r \in U(R)$. So x is clean and hence R is clean. $\square$

Corollary 4. *A directly finite ring $R \neq 0$ is local if and only if it is r -clean and 0 and 1 are the only idempotents in R .*

Theorem 5. *Let R be a commutative r -clean ring and each pair of idempotent in R be orthogonal. Then R is clean.*

Proof. By [1, Theorem 10], for commutative rings every regular ring is clean. So for each $x \in R$, we can write $x = e_1 + e_2 + u$, where $e_1, e_2 \in Id(R)$ and $u \in U(R)$. Now since e_1 and e_2 are orthogonal, $e = e_1 + e_2 \in Id(R)$. Hence $x = e + u$ is clean, which shows that R is clean. $\square$

Now, we give some properties of r -clean rings.

Theorem 6. *Let R be a ring, then $x \in R$ is r -clean if and only if $1 - x$ is r -clean.*

Proof. Let $x \in R$ be r -clean. Then write $x = r + e$, where $r \in Reg(R)$ and $e \in Id(R)$. Thus $1 - x = -r + (1 - e)$. But there exists $y \in R$ such that $ryr = r$. Hence $(-r)(-y)(-r) = -(ryr) = -r$ and since $-r \in Reg(R)$ and $1 - e \in Id(R)$, so $1 - x$ is r -clean.

Conversely, if $1 - x$ is r -clean, write $1 - x = r + e$, where $r \in Reg(R)$ and $e \in Id(R)$. Thus $x = -r + (1 - e)$, like previous part, $-r \in Reg(R)$ and $1 - e \in Id(R)$. Therefore x is r -clean. $\square$

Corollary 7. *Let R be a ring and $x \in J(R)$. Then x is r -clean.*

Proof. Let $x \in J(R)$. Then $1 - x \in U(R)$. So $1 - x \in Reg(R)$. Hence $1 - x$ is r -clean. Therefore by Theorem 6, x is r -clean. $\square$

Theorem 8. *Every factor ring of an r -clean ring is r -clean. In particular a homomorphic image of an r -clean ring is r -clean.*

Proof. Let R be r -clean and $I \triangleleft R$. Also let $\bar{x} = x + I \in \frac{R}{I}$. Since R is r -clean so we have $x = r + e$, where $r \in Reg(R)$ and $e \in Id(R)$. Thus $\bar{x} = \bar{r} + \bar{e}$. But there exists $y \in R$ such that $ryr = r$. Therefore $\bar{r}\bar{y}\bar{r} = \bar{r}$. So $\bar{r} \in Reg(R)$ and since $\bar{e} \in Id(\frac{R}{I})$, it follows that $\frac{R}{I}$ is r -clean. $\square$

Remark 9. *In general, inverse of above theorem may not be correct. For example, if p be a prime number, then $\frac{\mathbb{Z}}{p\mathbb{Z}} \cong \mathbb{Z}_p$ is r -clean, but $\mathbb{Z}$ is not r -clean.*

Theorem 10. *A direct product $R = \prod_{i \in I} R_i$ of rings $\{R_i\}_{i \in I}$ is r -clean if and only if so is each $\{R_i\}_{i \in I}$.*

Proof. One direction immediately follows from Theorem 8.

Conversely, let R_i be r -clean for each $i \in I$. Set $x = (x_i)_{i \in I} \in \prod_{i \in I} R_i$. For each i , write $x_i = r_i + e_i$, where $r_i \in Reg(R_i)$ and $e_i \in Id(R_i)$. Since $r_i \in Reg(R_i)$, there exists $y_i \in R_i$ such that $r_i y_i r_i = r_i$. Thus $x = (r_i)_{i \in I} + (e_i)_{i \in I}$, where $(r_i)_{i \in I} \in Reg(\prod_{i \in I} R_i)$ and $(e_i)_{i \in I} \in Id(\prod_{i \in I} R_i)$. Therefore $\prod_{i \in I} R_i$ is r -clean. $\square$

Lemma 11. *Let R be a commutative ring and $f = \sum_{i=0}^n a_i x^i \in R[x]$ be regular. Then a_0 is regular and a_i is nilpotent for each i .*

Proof. Since f is regular, thus there exists $g = \sum_{i=0}^m b_i x^i \in R[x]$ such that $fgf = f$. So $a_0 b_0 a_0 = a_0$. Therefore a_0 is regular. Now to end the proof, it is enough to show that for each prime ideal P of R ; every $a_i \in P$. Since P is prime, thus $\frac{R}{P}[x]$ is an integral domain. Define $\varphi : R[x] \rightarrow \frac{R}{P}[x]$ by $\varphi(\sum_{i=0}^k a_i x^i) = \sum_{i=0}^k (a_i + p)x^i$. Clearly φ is an epimorphism. But $\varphi(f)\varphi(g)\varphi(f) = \varphi(f)$, so $\deg(\varphi(f)\varphi(g)\varphi(f)) = \deg(\varphi(f))$. Thus $\deg(\varphi(f)) + \deg(\varphi(g)) + \deg(\varphi(f)) = \deg(\varphi(f))$. Therefore $\deg(\varphi(f)) + \deg(\varphi(g)) = 0$. So $\deg(\varphi(f)) = 0$. Thus $a_1 + P = \dots = a_n + P = P$, as required. $\square$

Theorem 12. *If R is a commutative ring, then $R[x]$ is not r -clean.*

Proof. We show that x is not r -clean in $R[x]$. Suppose that $x = r + e$, where $r \in \text{Reg}(R[x])$ and $e \in \text{Id}(R[x])$. Since $\text{Id}(R) = \text{Id}(R[x])$ and $x = r + e$, so $x - e$ is regular. Hence by pervious Lemma, 1 should be nilpotent, which is a contradiction. $\square$

Remark 13. *Even if R is a field, then $R[x]$ is not r -clean.*

Corollary 14. *If R is a commutative ring, then $R[x]$ is neither clean nor regular.*

Theorem 15. *Let R be a ring. Then the ring $R[[x]]$ is r -clean if and only if so is R .*

Proof. If $R[[x]]$ is r -clean, then by Theorem 8, $R \cong \frac{R[[x]]}{(x)}$ is r -clean. Conversely, suppose that R is r -clean. We know that $R[[x]] \cong \{(a_i) : a_i \in R, \text{ for each } i \geq 0\} = \prod_{i \geq 0} R$. So the result is clear by Theorem 10. $\square$

Theorem 16. *For every ring R , we have the following statements:*

- (1) *If e is an central idempotent element of R and eRe and $(1 - e)R(1 - e)$ are both r -clean, then so is R .*
- (2) *Let $e_1, \dots, e_n$ be orthogonal central idempotents with $e_1 + \dots + e_n = 1$. Then $e_i R e_i$ is r -clean for each i , if and only if R is r -clean.*
- (3) *If R is r -clean, then so is the matrix ring $M_n(R)$ for any $n \geq 1$.*
- (4) *If R is r -clean and M is a free R -module of rank n , then $\text{End}(M)$ is r -clean.*
- (5) *If A and B are rings and $M = {}_B M_A$ is a bimodule, the formal triangular matrix ring $T = \begin{pmatrix} A & 0 \\ M & B \end{pmatrix}$ is r -clean, then both A and B are r -clean.*
- (6) *Let for each integer $n \geq 2$, the ring T of all $n \times n$ lower(resp., upper) triangular matrices over R be r -clean. Then R is r -clean.*

Proof. We use $\bar{e}$ to denote $1 - e$ and apply the Pierce decomposition for the ring R , i.e.,

$$R = eRe \oplus eR\bar{e} \oplus \bar{e}Re \oplus \bar{e}R\bar{e}.$$

But idempotents in R are central, so $R = eRe \oplus \bar{e}R\bar{e} \cong \begin{pmatrix} eRe & 0 \\ 0 & \bar{e}R\bar{e} \end{pmatrix}$.

For each $A \in R$, write $A = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$, where a, b belong to eRe and $\bar{e}R\bar{e}$,

respectively. By our hypothesis a, b are r -clean. Thus $a = r_1 + e_1$, $b = r_2 + e_2$, where $r_1, r_2 \in \text{Reg}(R)$ and $e_1, e_2 \in \text{Id}(R)$. So

$$A = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} = \begin{pmatrix} r_1 + e_1 & 0 \\ 0 & r_2 + e_2 \end{pmatrix} = \begin{pmatrix} r_1 & 0 \\ 0 & r_2 \end{pmatrix} + \begin{pmatrix} e_1 & 0 \\ 0 & e_2 \end{pmatrix}.$$

But there exists $y_1, y_2 \in R$ such that $r_1 y_1 r_1 = r_1$, $r_2 y_2 r_2 = r_2$. Therefore

$$\begin{pmatrix} r_1 & 0 \\ 0 & r_2 \end{pmatrix} \begin{pmatrix} y_1 & 0 \\ 0 & y_2 \end{pmatrix} \begin{pmatrix} r_1 & 0 \\ 0 & r_2 \end{pmatrix} = \begin{pmatrix} r_1 y_1 r_1 & 0 \\ 0 & r_2 y_2 r_2 \end{pmatrix} = \begin{pmatrix} r_1 & 0 \\ 0 & r_2 \end{pmatrix}.$$

So $\begin{pmatrix} r_1 & 0 \\ 0 & r_2 \end{pmatrix} \in \text{Reg}(R)$, since $\begin{pmatrix} e_1 & 0 \\ 0 & e_2 \end{pmatrix} \in \text{Id}(R)$, it follows that R is r -clean.

On direction of (2) follows from (1) by induction. Conversely, let R be a r -clean ring and $e_1, \dots, e_n$ be orthogonal central idempotents with $e_1 + \dots + e_n = 1$. Then since $R = e_1 R e_1 \oplus \dots \oplus e_n R e_n$ then by Theorem 8, $e_i R e_i$ is r -clean for each i .

Also (3) follows from (2), and (4) follows from (3) and the fact that $\text{End}(R^n) \cong M_n(R)$.

For the proof of (5), let $T = \begin{pmatrix} A & 0 \\ M & B \end{pmatrix}$ be r -clean. Then for every $a \in A$, $b \in B$ and $m \in M$, write $\begin{pmatrix} a & 0 \\ m & b \end{pmatrix} = \begin{pmatrix} f_1 & 0 \\ f_2 & f_3 \end{pmatrix} + \begin{pmatrix} r_1 & 0 \\ r_2 & r_3 \end{pmatrix}$, where $\begin{pmatrix} f_1 & 0 \\ f_2 & f_3 \end{pmatrix} \in \text{Id}(T)$ and $\begin{pmatrix} r_1 & 0 \\ r_2 & r_3 \end{pmatrix} \in \text{Reg}(T)$. So $a = f_1 + r_1$ and $b = f_3 + r_3$. But there exists $\begin{pmatrix} y_1 & 0 \\ y_2 & y_3 \end{pmatrix}$ such that

$$\begin{pmatrix} r_1 & 0 \\ r_2 & r_3 \end{pmatrix} \begin{pmatrix} y_1 & 0 \\ y_2 & y_3 \end{pmatrix} \begin{pmatrix} r_1 & 0 \\ r_2 & r_3 \end{pmatrix} = \begin{pmatrix} r_1 & 0 \\ r_2 & r_3 \end{pmatrix}.$$

So

$$\begin{pmatrix} r_1 y_1 r_1 & 0 \\ r_2 y_1 r_1 + r_3 y_2 r_1 + r_3 y_3 r_2 & r_3 y_3 r_3 \end{pmatrix} = \begin{pmatrix} r_1 & 0 \\ r_2 & r_3 \end{pmatrix}.$$

Hence $r_1 \in \text{Reg}(A)$ and $r_3 \in \text{Reg}(B)$. It is easy to check that $f_1 \in \text{Id}(A)$ and $f_3 \in \text{Id}(B)$. Therefore a and b are r -clean. Hence both A and B are r -clean.

To end the proof, we can see (6) follows from (5). $\square$

Proposition 17. *Let R be a r -clean ring and e be a central idempotent in R . Then eRe is also r -clean.*

Proof. Since e is central, it follows that eRe is homomorphic image of R . Hence the result follows from Theorem 8. $\square$

Theorem 18. *Let R be a ring in which 2 is invertible. Then R is r -clean if and only if every element of R is the sum of a regular and a square root of 1.*

Proof. Suppose that R is r -clean and $x \in R$, then $\frac{x+1}{2} \in R$. Write $\frac{x+1}{2} = r + e$, where $r \in \text{Reg}(R)$ and $e \in \text{Id}(R)$. So $x = (2e - 1) + 2r$. But there exists $y \in R$ such that $ryr = r$. Thus $(r+r)\frac{y}{2}(r+r) = \frac{ryr}{2} + \frac{ryr}{2} + \frac{ryr}{2} + \frac{ryr}{2} = \frac{1}{2}(r+r+r+r) = 2r$. Thus $2r \in \text{Reg}(R)$ and since $(2e - 1)^2 = 1$, so x is a

sum of a regular and a square root of 1.

Conversely, if $x \in R$, then $2x - 1 = t + r$, where $t^2 = 1$ and $r \in \text{Reg}(R)$. Thus $x = \frac{t+1}{2} + \frac{r}{2}$. It is easy to check that $\frac{t+1}{2} \in \text{Id}(R)$. Now since $\frac{r}{2}(y + y)\frac{r}{2} = \frac{ryr}{4} + \frac{ryr}{4} = \frac{ryr}{2}$, it follows that $\frac{r}{2} \in \text{Reg}(R)$. Hence x is r -clean, which shows that R is r -clean. $\square$

If G is a group and R is a ring we denote the group ring over R by RG . If RG be r -clean, then R is r -clean by Theorem 8. But it seems to be difficult to characterize R and G for which RG is r -clean in general. In [4], [6] and [8] have given some rings and groups that RG is clean (so is r -clean). In following we will give some rings and groups that RG is r -clean.

Theorem 19. *Let R be a commutative semiperfect ring, G be a group and $(eRe)G$ be r -clean for each local idempotent e in R . Then RG is r -clean.*

Proof. Since R is semiperfect, so by [2, Theorem 27.6], R has a complete orthogonal set $e_1, \dots, e_n$ of idempotents with each e_iRe_i a local ring for each i . So e_i is a local idempotent for each i . Now by hypothesis, $(e_iRe_i)G$ is r -clean. Since $(e_iRe_i)G \cong e_i(RG)e_i$ for each i , it follows that $e_i(RG)e_i$ is r -clean. Hence RG is r -clean by Theorem 16 (2). $\square$

Theorem 20. *Let R be a ring which 2 is invertible and $G = \{1, g\}$ be a group. Then RG is r -clean if and only if R is r -clean.*

Proof. One direction is trivial.

Conversely, if R is r -clean, then since 2 is invertible by [6, Proposition 3], $RG \cong R \times R$. Hence RG is r -clean by Theorem 10. $\square$

REFERENCES

- [1] D. D. Anderson, V. P. Camillo, *Commutative rings whose elements are a sum of a unit and an idempotent*, Comm. Algebra 30 (7) (2002), 3327–3336.
- [2] F.W. Anderson, K.R. Fuller, *Rings and Categories of Modules*, Second edition, Graduate Texts in Mathematics, 13, Springer-Verlag, New York, 1992.
- [3] V. P. Camillo, H. -P. Yu, *Exchange rings, unit and idempotents*, Comm. Algebra 22 (12)(1994), 4737–4749.
- [4] J. Chen, W. K. Nicholson, Y. Zhou, *Group ring in which every element is uniquely the sum of a unit and an idempotent*, J. Algebra 306 (2)(2006), 453–460.
- [5] K.R. Goodearl, von Neumann Regular Rings, 2nd ed., Robert E. Krieger Publishing Co. Inc., Malabar, FL, 1991.
- [6] J. Han, W. K. Nicholson, *Extensions of clean rings*, Comm. Algebra 29 (6) (2001), 2589–2595.
- [7] D. Handelman, *Perspectivity and cancellation in regular rings*, J. Algebra 48 (1977), 1–16.
- [8] W. Wm. McGovern, *A characterization of commutative clean rings*, Int. J. Math. Game Theory Algebra 15(4) (2006), 403–413.
- [9] W. K. Nicholson, *Lifting idempotent and exchange rings*, Trans. Amer. Math. Soc. 229 (1977), 269–278.
- [10] W. K. Nicholson, K. Varadarajan, *Countable liner transformations are clean*, Proc. Amer. Math. Soc. 126 (1998), 61–64.
- [11] W. K. Nicholson, Y. Zhou, *Rings in which elements are uniquely the sum of an idempotent and a unit*, Glasg. Math. J. 46 (2) (2004), 227–236.

DEPARTMENT OF MATHEMATICS, SEMNAN UNIVERSITY, SEMNAN, IRAN,
E-mail address: <ebrahimnasibi@yahoo.com>, <enasibi@gmail.com>